



E1048

JACC March 27, 2012

Volume 59, Issue 13



Heart Failure

OBSTRUCTIVE SLEEP APNEA AND ASTHMA CO-MORBIDITIES ARE INDEPENDENT PREDICTORS OF NEW HEART FAILURE AND ATRIAL FIBRILLATION IN PATIENTS WITH DIABETES MELLITUS AND PRESERVED EJECTION FRACTION

ACC Moderated Poster Contributions

McCormick Place South, Hall A

Monday, March 26, 2012, 11:00 a.m.-Noon

Session Title: Heart Failure Complicated by Anemia or Diabetes Mellitus

Abstract Category: 14. Heart Failure: Clinical

Presentation Number: 1230-619

Authors: *Sinziana Seicean, Kazuaki Negishi, Tomoko Negishi, Teerapat Yingchoncharoen, Conrad Gibby, Thomas Cook, Andreea Seicean, Xuebei An, Thomas Marwick, Heart and Vascular Institute, Cleveland Clinic, Cleveland, OH, USA, Case Western Reserve University, Cleveland, OH, USA*

Background: Heart failure (HF) and atrial fibrillation (AF) are twin epidemics with poor prognosis in pts with type 2 diabetes mellitus (T2DM). Obstructive sleep apnea (OSA) is an important correlate of T2DM, but its role in HF+AF is unclear, as is the role of underlying lung disease. We sought to evaluate clinical predictors of new HF and AF in T2DM.

Methods: 914 consecutive T2DM pts (age 56 ± 11 y) without prior HF or AF and with a negative stress echo were enrolled in an observational study, after exclusion of prior cardiac surgery, > mild valvular heart disease, or ejection fraction < 50%. Demographics, clinical assessments based on standards of diabetes care, co-morbidities, and treatment with insulin, diuretics, beta-blockers, statins, ace-inhibitors and aspirin were collected at the time of stress echo. Pts were followed for up to 8 y for HF+AF, and best predictors of endpoints were sought by using nested Cox proportional hazard models.

Results: 22 new HF and 25 new AF developed during a mean time of 4.8 y. OSA and asthma were associated with outcome in univariate and all adjusted models (Table), independent of socio-economic class, diabetes care and clinical predictors.

Conclusion: OSA and asthma are associated with HF+AF in T2DM.

Table: Impact of Baseline OSA and Asthma on HF+AF in Cox Models:

	OSA		Asthma	
	Hazard Ratio (95% CI)	p-value	Hazard Ratio (95% CI)	p-value
Univariate	2.3 (1.01-5.04)	0.046	2.3 (1.1, 4.8)	0.037
Multivariate (OSA, Asthma) with:				
Age, gender	2.6 (1.1-5.9)	0.025	2.6 (1.2-5.6)	0.019
Age, gender, race, poverty	2.6 (1.2-6.1)	0.022	2.5 (1.2-5.6)	0.021
Age, gender, ejection fraction, left atrial volume, nonoptimal systolic or diastolic pressure at baseline echo	2.5 (1.1-5.6)	0.034	2.5 (1.1-5.6)	0.022
Age, gender, hypertension, obesity, CAD, resistant hypertension, depression, anxiety, neuropathy, nephropathy	2.5 (1.1-5.7)	0.035	2.5 (1.1-6.0)	0.021
Age, gender, family history, smoking, hypertension, CAD	2.4 (1.04-5.5)	0.039	2.7 (1.2-6.0)	0.013
Age, gender, treatment	2.4 (1.03-5.5)	0.042	2.5 (1.1-5.6)	0.023
Age, Gender, nonoptimal control of diabetes, HDL, triglyceride and creatinine (2011 Diabetes Standards of Medical Care Guidelines)	2.6 (1.1-5.9)	0.0247	2.6 (1.2-5.7)	0.018